

indebted to Prof. A. Lakes and Engineer H. C. Beckwith of the U. S. Navy, who found the first remains in Colorado near the locality of the gigantic *Atlantosaurus montanus*, and in essentially the same horizon.

Yale College, New Haven, Nov. 15th, 1877.

ART. LIII.—Notice of New Dinosaurian Reptiles from the Jurassic formation; by Professor O. C. MARSH.

THE gigantic Dinosaur, *Atlantosaurus montanus*, described by the writer in the July number of this Journal,\* proves to belong to a lower horizon than at first supposed, and is really from the upper Jurassic. Additional remains of the type specimen, moreover, throw considerable light on the structure of this largest of land animals, and indicate that it is the representative of a distinct family, which may be called *Atlantosauridae*.

In the type genus, *Atlantosaurus*, one of the most important characters is the pneumaticity of the vertebræ, as mentioned in the original description. Another noteworthy feature is the absence on the femur of a third trochanter. The shaft of the bone is somewhat thickened at the point where this process should be, but the trochanter is wanting. The size of the original specimen of *A. montanus* may be estimated from the femur, which was about seven feet in length. If the animal had the proportions of a Crocodile, it was at least eighty feet long.

*Apatosaurus ajax*, gen. et sp. nov.†

Another gigantic Dinosaur, allied to the above, and of scarcely less interest, is represented in the Yale Museum by a nearly complete skeleton in excellent preservation. It is from the Jurassic beds in the Eastern foot hills of the Rocky Mountains, but from a somewhat lower horizon than the type of *Atlantosaurus*.

The cervical vertebræ are strongly opisthocœlous, and are rendered comparatively light by large pneumatic cavities in the centra. The anterior dorsals have similar characters. The posterior lumbar have the articular faces very nearly flat, and transverse. The sacral vertebræ are more solid, and have their transverse processes nearer the middle of the centra than in *Atlantosaurus*. The anterior caudals are biconcave, and their

\* Vol. xiv, p. 87, 1877. The name *Titanosaurus* was first given, but, being preoccupied, may be replaced by *Atlantosaurus*.

† The principal characters of this genus and its nearest allies were given by the writer in a paper before the National Academy of Science, at the meeting in New York, October 25th, 1877.

interior structure is cancellous. The chevron bones differ from those of most known Dinosaurs in having the superior articular ends of the rami not united, but separated from each other, as in the Mosasauria with free hæmapophyses.

Some of the dimensions of this skeleton are as follows:

Length of centrum of anterior dorsal vertebra .....	220 <sup>mm</sup>
Transverse diameter of anterior face .....	310
Vertical diameter .....	200
Amount of convexity .....	90
Length of centrum of lumbar vertebra .....	240
Transverse diameter of anterior face .....	410
Vertical diameter .....	270
Length of median sacral vertebra .....	250
Expanse of its transverse processes .....	760
Length of centrum of median caudal .....	190
Length of posterior caudal .....	185

This animal must have been between fifty and sixty feet in length, and more than thirty in height when erect.

*Apatosaurus grandis*, sp. nov.

Another huge Dinosaur, apparently of the same genus, but of smaller size, is represented in the Yale Museum by the more important parts of a skeleton, in remarkable preservation. In this specimen the cervical vertebræ have the walls of the centra very thin. The caudals preserved are elongated and slender, indicating a long tail. The femur is comparatively short, and without a third trochanter. The great trochanter is much lower than the head of the femur, and continuous with it. The metapodial bones indicate a foot of medium length.

The following measurements indicate the size of the reptile:

Length of femur .....	1050 <sup>mm</sup>
Transverse diameter of proximal end .....	340
Transverse diameter of distal end .....	290
Length of posterior caudal vertebra .....	145
Vertical diameter of anterior articular face .....	110
Transverse diameter .....	115

The known remains of this species are from the same geological horizon as those above described. They indicate an animal at least thirty feet in length.

*Allosaurus fragilis*, gen. et sp. nov.

This genus may be distinguished from any known Dinosaurs by the vertebræ, which are peculiarly modified to ensure lightness. Although apparently not pneumatic, they have the weight of the centra greatly reduced by deep excavations in the sides. Some of them have the centra hour-glass in form,

the middle part being so diminished as to greatly reduce the strength. The vertebræ preserved are biconcave, with shallow cavities. The feet bones referred to this species are very slender. A lumbar vertebra has its centrum 105 mm. in length, and 89 in least transverse diameter. An anterior caudal, 35 mm. long, has its centrum so much constricted that its least transverse diameter is 38 mm., while its anterior face is 90 mm. in transverse diameter.

The animal indicated by the remains preserved was from fifteen to twenty feet in length. All the known specimens are from the upper Jurassic of Colorado.

*Nanosaurus rex*, sp. nov.

A diminutive Dinosaur, about as large as a fox, is indicated by some remains in good preservation, the most characteristic of which is a nearly perfect femur. In this bone, the great trochanter is prominent, and the third trochanter especially so. There is a well developed fibular ridge, directed outward and backward. The cavity in this bone is unusually large, and the walls are smooth. This femur agrees so nearly with that of the type of *Nanosaurus*, that the present species may be provisionally referred to that genus.

The dimensions of this bone are as follows:

Length of femur .....	100· mm
Distance from head to middle of third trochanter ..	30·
Transverse diameter of distal end .....	21·
Greatest antero-posterior diameter .....	18·
Least transverse diameter of shaft .....	11·
Diameter across third trochanter .....	15·

The known remains of this reptile are from the upper Jurassic of Colorado.

The specimens described in the present articles are deposited in the Peabody Museum of Yale College. They are all from essentially the same geological horizon, which I find to be upper Jurassic. The deposits which contained them may be called the *Atlantosaurus* beds, from their most characteristic fossils, the huge Dinosaurs of that genus.

Yale College, New Haven, November, 1877.

## INDEX TO VOLUME XIV.\*

### A

- Abbott, H. L.*, artificial tremors through the earth's crust, 509.
- Academy*, National, publications of, 167. October meeting, 511.. Nat. Sci. Phil., Journal of, 78. of Sciences, St. Petersburg, 167. Wisconsin, transactions of, 78.
- Acid*, phyllic, from leaves, 483. rosolic, from cresol and phenol, 414. salicylic, method of producing, 66. tartaric, from pyruvic, 310.
- Acids*, constitution of unsaturated dibasic, 413. complex inorganic, *Gibbs*, 61.
- Agassiz, A.*, N. American star-fishes, 73. zoological notices, 500.
- Airy, G.*, sun's distance, 501.
- Allen, J. A.*, influence of physical conditions in the genesis of species, 161. North American rodentia, not., 422.
- Allen, O. D.*, hatchettolite and samarskite, 128.
- Amylene* from amyl iodide, 412.
- Anthropology*, *Galton*, 265.
- Archæologists*, caution to, 333.
- Archæology*, Peabody Museum of, Report, 246.
- Armsby, H. P.*, absorption of bases by the soil, 25.
- Association*, American, 76. Nashville meeting, 328. Marsh's address, 337. British, Plymouth, 334. Galton's address, 265.
- Astronomical observations*, Harvard College, 75. Cincinnati, 246. Washington, 74.
- Atterberg*, the terpenes of Swedish wood tar, 412.
- Auria*, conversion of into rosaniline, 310.
- Autography*, practical use of, *Sars*, 277.
- B**
- Baker, J. G.*, Iridaceæ, 428.
- Barker, G. F.*, chemical abstracts, 64, 148, 309, 411, 481.
- Barnard, C.*, Light, 419.
- Bean, T. H.*, two new species of fishes, 470.
- Becquerel, H.*, rotatory polarization, 417.
- Bennett, A. W.*, rapid growth, 243.
- Bermudas*, fishes of, *Goode*, 289.
- Berthelot*, effect of pressure on chemical action, 64.
- Bolton, H. C.*, organic acids in examination of minerals, 495.
- Börnstein*, influence of light upon electrical resistance of metals, 152.
- BOTANY—**
- Algae*, North American, 72.
- "*Artichokes*," native, 428.
- Athamanta Chinensis*, 160.
- Flora Brasiliensis*, 427.
- Fungi*, diseases caused by, 426.
- Growth*, abnormal, in an apple-tree, *Meehan*, 243. rapid, 243.
- Growth-rings* in exogens, *Warring*, 394.
- Impatiens*, cleistogamy in, 497.
- Lichens*, reproductive organs of, 72.
- Megarrhiza*, germination of, *Gray*, 21.
- New Jersey*, catalogue of plants, 498.
- Nomenclature*, *Gray*, 158.
- Orchis rotundifolia*, *Gray*, 72.
- Pithophoraceæ*, 71.
- Rocky Mountains*, *Hooker*, 505.
- Wild flowers of America*, 497.
- See also under **GEOLOGY**.
- Bottomley, J. T.*, Dynamics, 168.
- Bougarel*, phyllic acid, extracted from leaves, 483.
- Bourgoin*, action of bromine upon pyrotartaric acid, 150.
- Boutlerow*, isobutylene, 66.
- Brohnsiesig, G. C. W.*, Year book of botanical literature, 160.
- Bromine*, action of, upon pyrotartaric acid, 150.
- Burnham, S. W.*, double-star discoveries, 31.
- Bussey Institution*, Bulletin of, 168.
- C**
- Capillarity*, Gauss's theory of, 152.
- Carbonates and oxalates*, 482.
- Carnelley*, determination of high melting points, 65.
- Caton, J. D.*, Antelope and deer of America, 426.
- Cavern exploration* in Devonshire, *Pengelly*, 299.

\* The Index contains the general heads BOTANY, GEOLOGY, MINERALOGY, ZOOLOGY, and under each the titles of Articles referring thereto are mentioned.